

REPORT DOCUMENTATION PAGE				<i>Form Approved</i> OMB No. 0704-0188		
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Service Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.</p>						
1. REPORT DATE (DD-MM-YYYY) 20/2/2010		2. REPORT TYPE AFOSR Final Progress Report		3. DATES COVERED (From - To) 1/1/2007 to 30/11/2009		
4. TITLE AND SUBTITLE Using Attosecond Technology to Image Molecular Orbitals				5a. CONTRACT NUMBER FA9550-07-1-0068		
				5b. GRANT NUMBER AFOSR		
				5c. PROGRAM ELEMENT NUMBER 		
6. AUTHOR(S) Dr. Paul Corkum				5d. PROJECT NUMBER FIATA06319B005		
				5e. TASK NUMBER 		
				5f. WORK UNIT NUMBER 		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) National Research Council of Canada 100 Sussex Drive Ottawa, ON, K1A 0R6				8. PERFORMING ORGANIZATION REPORT NUMBER ATTOSECOND		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) National Research Council of Canada 100 Sussex Drive Ottawa, ON, K1A 0R6				10. SPONSOR/MONITOR'S ACRONYM(S) NRC		
				11. SPONSOR/MONITOR'S REPORT NUMBER(S) AFRL-OSR-VA-TR-2012-0603		
12. DISTRIBUTION/AVAILABILITY STATEMENT Distribution A : Approved for Public Release						
13. SUPPLEMENTARY NOTES None						
14. ABSTRACT <p>Attosecond hole dynamics: While studying the spectrum of the high harmonics produced from CO₂, we discovered that multiple orbitals can ionize at the same time. This was an unexpected result suggesting that tunneling often launches attosecond dynamics. This research, including an image of the wave packet motion, was published in Nature [2].</p> <p>Imaging orbitals via tunneling: To gain additional insight into the ionization of lower orbitals, we measured the ionization probability as a function of direction. Using H₂ (because of its simplicity) we showed that the tunnel ionization rate essentially follows the orbital structure (Physical Review Letters [3]). Then, we turned to HCl and measured the relative probability of tunnel ionizing the HOMO and the HOMO-1 orbital (Science [1]). Finally we developed an electron tomographic imaging technique [6].</p>						
15. SUBJECT TERMS ATTOSECOND, Molecular Orbitals, Imaging, High harmonics, tunneling, ionization						
16. SECURITY CLASSIFICATION OF: a. REPORT b. ABSTRACT c. THIS PAGE			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON 19b. TELEPHONE NUMBER (Include area code)	

Annual Accomplishments:

Our objective for the whole AFOSR proposal was to image molecular orbitals with pi symmetry. Below we outline our accomplishments during 2009:

Attosecond hole dynamics: While studying the spectrum of the high harmonics produced from CO₂ we uncovered that multiple orbitals were ionizing at the same time when the molecule was viewed from some directions. This was an unexpected and exciting result because it implied that a hole wave packet was launched in the molecule. We measured this wave packet which, in CO₂, moved with a natural time scale of ~ 1 fs. This research, suggesting that tunneling often launches previously unsuspected attosecond dynamics in molecules, was published in *Nature* as a full article [2].

Imaging orbitals via tunneling: To gain additional insight into the ionization of lower orbitals we measured the ionization probability as a function of direction in two molecules. We named this new orbital imaging method a “molecular STM” because of its parallel with a conventional “scanning tunneling microscope”. Using H₂ (because of its simplicity) in 2008 we showed that the tunnel ionization rate in 800 nm light varies as a function of angle essentially following the orbital structure. This work was published during 2009 in *Physical Review Letters* [3]. Exploiting this fact, we turned to HCl, and Confirmed that not only the HOMO, but also the HOMO-1 ionized. This work was published in *Science* [1]. Finally we improved experimental methods for measuring tunneled electrons by developing an electron tomographic imaging technique [6].

Infrared experiments: Tunneling is a better approximation for ionization if done with infrared radiation. Orbital tomography is also preferentially performed with infrared radiation because the recollision electron wave length is shorter. In a paper published in *Physical Review Letters* we studied the wavelength scaling of high harmonic generation using infrared radiation [4]. As the year drew to an end we extended our infrared experiments to more complex molecules [5].

Publications during 2009:

1. H. Akagi, T. Otobe, A. Staudte, A. Shiner, F. Turner, R. Dörner, D.M. Villeneuve and P.B. Corkum, “Direct Observation of Tunneling from HOMO-1 in HCl”, *Science*, **325**, 1364 (2009).
2. O. Smirnova, Y. Mairesse, S. Patchkovskii, N. Dudovich, D. M. Villeneuve, P. B. Corkum, and M. Y. Ivanov, “Probing Inner Orbitals and Electron Tunneling in Molecules with High Harmonic Interferometer”, *Nature*, **460**, 972 (2009).
3. A. Staudte, S. Patchkovskii, D. Pavicic, H. Akagi, O. Smirnova, D. Zeidler, M. Meckel, D.M. Villeneuve, R. Dörner, M. Ivanov and P.B. Corkum, “Angular Tunneling Ionization Probability of H₂” *Phys. Rev. Lett.* **102**, 033004 (2009).

4. A. D. Shiner, C. Trallero-Herrero, N. Kajumba, H.-C. Bandulet, D. Comtois, F. Legare, M. Gigeure, J-C. Kieffer, P. B. Corkum, and D. M. Villeneuve, "Wavelength Scaling of High Harmonic Generation Efficiency", *Phys Rev Lett.* **103**, 073902 (2009).
5. C. Trallero-Herrero, B. E. Schmidt, A. D. Shiner, Philippe Lassonde, Eric Bisson, J-C. Kieffer, P. B. Corkum, D. M. Villeneuve, F. Legare, "High harmonic generation in Ethylene with infrared pulses", *Chemical Physics*, **366**, 33 (2009).
6. C. Smeenk, L. Arissian, A Staudte, D. M. Villeneuve, and P. B. Corkum, "Tomographic imaging of charged particles", *J. Phys. B.* **42**, 185402 (2009).

Changes to objectives: None

New program manager:

Extensions: None:

Final report: